




Nigerian Journal of Soil Science

Journal homepage: www.soilsjournalnigeria.com

Ammonium Sorption Characteristics of Soils of Coastal Plain Sand in Owerri West Imo State

¹Osisi A.F., ²Azu, D. E. O., ³Agim, L.C. and ⁴Chukwu, E.D^{1,3&4}Department of Soil Science and Technology, Federal University of Technology, Owerri, Imo State²Department of Horticulture and Landscape Technology, Akanulbiam Federal Polytechnic, Unwana, Ebonyi State

ARTICLE INFO

Article history:

Received April 27, 2023

Received in revised form May 21, 2023

Accepted June 9, 2023

Available online June 17, 2023

Keywords:

Ammonium

Coastal Plain Sand

Determination

Owerri

Sorption

Corresponding Author's E-mail Address:

donblessed01@gmail.com +234038663924

ISSN– Online 2736-1411

Print 2736-142X

© Publishing Realtime. All rights reserved.

ABSTRACT

Information on the ammonium sorption capacity of soils is very crucial in the development of sound ammonium fertilization and management. This study was therefore carried out to determine the ammonium sorption characteristics of soils of coastal plain sand formation in Owerri West LGA of Imo State. Samples were collected from three locations (FUTO, Ihiagwa and Umuokanne), at three depths (0-15, 15-30, 30-45 cm), giving a total of nine observational units. The experimental design was a 3 x 1 factorial in Randomized Complete Block Design (RCBD), the parent material and soil depth was the factors under consideration. Statistical analysis did not show significant ($P > 0.05$) variation in soil physical and chemical properties with location. However, most of the properties increased with depth. Data obtained from the analysis were statistically analyzed using Analysis of Variance (ANOVA) and means separated using Lsd(0.05). Correlations between soil routine properties and ammonium sorption parameters were done using SPSS (Version 16.0). Soil Ammonium sorption properties of the soils showed variability with location and depth. FUTO had the highest capacity to absorb ammonium (445mg/kg). For all locations, maximum adsorption capacity increased with depth. However, the affinity coefficient was greatest in soil of Ihiagwa (3.25mg/kg) and this increased with depth for all soils. The adsorption maximum and affinity coefficient correlated positively with clay and negatively with sand and organic matter. Langmuir and Freundlich isotherms were applied to test the sorption data. Comparing the fitness of the ammonium sorption models using the R^2 values, it was evident that Freundlich Model better described and predicated the ammonium sorption parameters and is therefore recommended for the study of soils of these locations.

1.0 Introduction

Ammonium in soil serves not only as a major source of Nitrogen for plant growth, but also as an important product or reactant in the Nitrogen transformation processes in soil (Brady and Weil, 2008). Ammonium is also the main form in which fertilizer Nitrogen is applied in agriculture. The composition of nitrogen fertilizers is dominated by ammonium nitrate, which provides half of the nitrogen in the ammonium form and half in the nitrate form (Zanget *et al.*, 2011). Ammonium is a prevalent Nitrogen subsurface pollutant, which emerges from different sources, such as landfill sites, industrial polluted areas, sewage systems, wastewater, and fertilizers (Ranjbar and Jalali, 2013). The concentrations of ammonium might exceed the standard

limits of disposal to water bodies (Zhu *et al.*, 2011), with high concentrations hindering the decomposition of organic matter via microorganisms (Ranjbar and Jalali, 2013). The three major routes of NH_4^+ removal in the ecosystems are mainly nitrification, assimilation into microbial cells, and adsorption onto substrate media (Sun *et al.*, 2005; Tyrrel *et al.*, 2002).

Ammonium adsorption or sorption involves the attraction or adhesion of ammonium to soil colloidal particle thereby preventing its release to crops (Brady and Weil, 2008). With the increased use of ammonium fertilizers, a study of the absorption, retention and release of NH_4^+ by soil is gaining considerable importance (Sutton *et al.*, 2011a). Adequate understanding of the ammonium sorption char-

acteristics of soils provides useful index for soil ammonium nitrogen management as this will promote efficient ammonium nitrogen utilization by crops to maximize yield. However, this information is lacking in the coastal plain sand of Owerri. In order to bridge this information gap, this study was embarked upon.

Materials and Method

2.1 Study Area

This study was carried out in three locations of Owerri West LGA, Imo State (Ihiagwa, Umuokanne and FUTO Campus). Ihiagwa lies between Latitude 5°40' N and Longitude 7°12' E, Umuokanne lies between 5°25' N and Longitude 6°58' E and FUTO lies between Latitude 5°29' N and Longitude 7°02' E. The rainfall of the area ranged from 1950mm to 2250mm per annum with average mean annual temperature of 27°C -29°C. (NIMET 2008). The dominated soil type is ultisols derived from Coastal plain sand and characterized by low Cation exchange Capacity, low organic matter, low pH and are highly weathered soils (Onweremmadu and Ufot, 2011).

2.2. Soil Sample Collection

Soil samples were collected from the identified locations (Ihiagwa, Umuokanne and FUTO Campus) at three depths (0-15cm and 15-30 and 30 -45cm) using soil auger. The samples were analyzed in the laboratory for routine analysis according to standard methods: Particle size distribution using hydrometer method according to the procedure of Gee and Or (2002), Bulk density (BD), according to Grossman and Reinsch (2002), soil pH was determined in soil to water and soil to CaCl_2 at a ratio 1:2 soil water and soil CaCl_2 respectively using glass electrode pH meter (Udo *et al.*; 2009). Organic carbon was determined by the wet oxidation method according to Pansu and Gautheyrou (2006) and converted to organic matter by multiplying by 1.792. The total nitrogen determination was done by the macro Kjeldahl digestion method (Simmons *et al.*, 1994). Available P was determined using the Bray II method of Bray as described by Udo *et al.*; (2009). Exchangeable acidity was determined by the nickel extraction procedure as described by Udo *et al.*, (2009). Exchangeable basic cations (K^+ , Ca^{2+} , Mg^{2+} , Na^+) were determined by the ammonium acetate method (Carter and Gregorich, 2008). Ca and Mg in the extract were determined using the atomic absorption spectrophotometer, while K and Na were determined using the flame photometer. Effective cation exchange capacity (ECEC) was obtained by summation of all the exchangeable cations and exchangeable acidity as described by Udo, *et al.*; (2009). The base saturation was obtained mathematically with

$$\text{BS (\%)} = \frac{\text{Total cation} \times 100}{\text{ECEC}} \quad 1$$

2.3 Ammonium Sorption Studies

Ammonium sorption study was conducted using Batch studies as described by Fox and Kamprath (1970). This briefly involves weighing 5g fine earth soil particle into a centrifuge tube and adding 25ml of NH_4^+ ppm solution containing graded concentrations of 0, 10, 20, 30, 40, and 50 g kg^{-1} of NH_4^+ ppm. The tube was shaken on a rotary shaker GFL 3040 with a speed of 14 minutes at 200rpm for 1 hr. After the equilibration time, the tube and its contents was centrifuged at 3500 x g for 30 mins and the clear supernatant decanted into the 50ml glass. Aliquots of the

supernatant were taken and color developed using a Nessler reagent.

The percent contribution of ammonium absorbed (%R) and ammonium adsorption capacity (q) was calculated using the following expression:

$$\%R = \frac{C_0 - C_e}{C_0} \times 100 \quad \text{.....(1)}$$

$$q_v = \frac{(C_0 - C_e)}{m} \quad \text{.....(2)}$$

Where C_0 - initial concentration of ammonium in the solution (Mg/L)

C_e - final ammonium concentration in the solution (Mg/L)

V - volume of the solution (mL)

M - amount of sorbent on the dry basis (g)

2.4 Statistical Analysis

Generated data was analyzed using analysis of variance (ANOVA). Significant means were separated using least significant difference (LSD) at 5% probability level. Correlation between soil properties and ammonium sorption parameters were determined using SPSS (16.0 version).

3.0 Results and Discussion

3.1 Physical and chemical properties of the soils used for the study

The physical and chemical properties of the soils used for the study is presented in tables 1 and 2. According to table 1, there is obvious variability in the textural properties of the soils among the different locations. FUTO had a textural class of loamy sand with sand occurring at above 75% and clay ranging from 6.96 to 8.96%. While sand decreased with depth, clay was found to be proportional to depth. According to Azuet *al.*, (2018), direct proportionality exists between clay content and soil depth in most weathered tropical soils of southeastern Nigeria. Ihiagwa and Umuokanne had similar textural class of sandy loam.

Bulk density and moisture content were low, while the total porosity was high. Sandy particle with their large pore spaces are prone high water drainage and low water and nutrient retention abilities (Brady and Weil, 2008). The clay component of the soil is very important in the determination of soil fertility properties, especially with regards to the potential of soils to retain or adsorb nutrients Azu *et al.*, (2018).

The chemical properties of the different soils did not show significant variation among soils of contracting locations, (Table 2). All of the different soil samples were slightly acidic for all depths with the pH generally above 6. However, soil depth did not cause obvious variability on pH. Organic matter was very high for all locations and sampling depths. Ihiagwa had the highest value of organic matter (9.31%), while FUTO had the least value (8.56%). For all locations, organic matter decreased with depth. Farming systems involving the addition of organic manure, planting of cover crops may have influenced this result. Onyegbule *et al.*, (2019), in their study of organic matter distribution reported decrease in organic matter concentration of soils down the soil profile.

Total nitrogen was high in all location and also showed a depth wise decline. Surface accumulation of organic mat-

Land Use	Sand %	Silt %	Clay %	S/C Ratio	Textural Class	Bd g/cm3	Tp %	Mc %
	Futo							
0-15	85.76	7.28	6.96	1.05	LS	1.05	42.00	17.54
15-30	83.76	8.28	7.96	1.04	LS	1.54	60.40	23.31
30-45	79.76	11.28	8.96	1.26	LS	1.42	46.40	15.32
Mean	83.09	8.94	7.96	1.12		1.34	49.6	18.7
CV	11.22	144.7	12.96	1.25		2.99	186.1	90.91
CV Ranking	*	***	*	*		*	***	**
	Ihiagwa							
0-15	79.76	9.28	10.96	0.85	SL	1.54	41.90	13.31
15-30	67.76	19.28	12.96	1.49	SL	1.05	42.00	19.06
30-45	69.76	15.28	14.96	1.02	SL	1.54	41.90	13.30
Mean	72.43	14.61	12.94	1.12		1.38	41.93	15.22
CV	57.29	173.3	30.91	10.27		5.79	0.01	72.47
CV Ranking	**	***	**	*		*	*	**
	Umuokanne							
0-15	77.76	11.28	10.96	1.01	SL	1.05	42.00	17.50
15-30	79.76	12.28	7.96	1.53	LS	1.54	60.40	23.29
30-45	65.76	22.28	11.96	1.86	SL	1.42	46.40	14.32
Mean	74.43	15.28	10.13	1.47		1.34	49.6	18.37
CV	77.02	242.2	43.04	13.05		4.72	186.1	112.6
CV Ranking	*	***	*	*		*	**	**

LAND USE	pH(H ₂ O)	Oc%	OM%	TN %	Av. P (PPM)	Ca ²⁺	Mg ²⁺	Na+	K+	Al ^{3++H}	Al ³⁺	TEA	TEB	%BS	ECEC cmol/kg
SITE1							FUTO								
0-15	6.98	5.79	9.98	0.32	20.44	1.37	0.01	0.006	0.008	0.334	-	0.334	1.39	80.67	1.73
15-30	7.01	4.65	8.02	0.15	9.52	0.11	0.11	0.017	0.007	1.336	-	1.244	0.24	15.24	1.58
30-45	6.98	4.45	7.67	0.116	18.62	0.27	0.15	0.006	0.005	1.503	-	0.431	0.43	22.29	1.93
Mean	6.99	4.96	8.56		16.19	0.58	0.12	0.009	0.007	1.06		0.67	0.69	39.4	1.75
CV	0.005	10.56	18.11		211.4	81.03	9.17	0.46	0.04	37.74		37.31	56.52	3273.9	1.89
CV Rank-ing LSD	*	*	*		**	*	*	*	*	*		*	*	***	*
SITE2	Ihiagwa														
0-15	6.42	5.79	9.98	0.355	19.11	0.93	0.13	0.008	0.007	0.501	-	1.075	1.08	68.21	1.58
15-30	6.22	5.29	9.12	0.256	22.61	0.19	0.43	0.027	0.015	0.668	0.48	0.662	0.66	36.57	1.81
30-45	6.13	5.13	8.84	0.197	16.17	0.25	0.07	0.01	0.009	0.083	-	0.319	0.32	79.35	0.4
Mean	6.26	5.4	9.31		29.29	0.46	0.21	0.015	0.01	0.42		0.69	0.69	61.38	1.26
CV	0.46	2.41	3.87		53.76	36.96	17.62	0.73	0.18	21.59		20.72	20.72	802.5	45.24
CV Rank-ing LSD	*	*	*		**	**	*	*	*	*		*	*	***	**
SITE3	Umuokanne														
0-15	6.06	5.25	9.05	0.216	19.6	0.51	0.19	0.01	0.014	1.67	-	0.724	0.72	30.24	2.39
15-30	6.31	4.73	8.15	0.149	25.48	0.13	0.33	0.008	0.006	1.336	-	0.474	0.47	26.18	1.81
30-45	6.57	4.99	8.60	0.132	20.72	0.07	0.08	0.009	0.016	0.668	-	0.175	0.18	20.76	0.84
Mean	6.31	4.99	8.6		21.93	0.24	0.2	0.009	0.012	1.22		0.46	1.37	25.73	1.68
CV	1.11	1.6	3.63		44.41	25	7.85	0.01	0.38	21.31		16.74	96.64	87.91	36.31
CV Rank-ing LSD	*	*	*		**	**	*	*	*	**		*	***	***	**

3.2 Ammonium Sorption Parameters using Langmuir and Freundlich Models

The ammonium sorption characteristics of the soils using

Freundlich and Langmuir's equations are shown in table 3.

According to this Table, the ammonium sorption maximum was highest in soils of Ihiagwa for Langmuir model

Table 3.. Ammonium sorption characteristics of the soil profiles using Langmuir and Freundlich equations at different locations and soil depth

Location	Depth	Langmuir				Freundlich			
		b	k	MBC	R ²	a	n	MBC	R ²
	(cm)			(mg/kf)				(mg/kg)	
FUTO	0 - 15	112	7.4	829	0.185	250	1.4	350	0.849
	15 - 30	170	5.9	1003	0.180	252	1.4	353	0.873
	30 - 45	278	1.8	500	0.505	220	0.08	176	0.800
Mean		187	5.03	777.3	0.290	240.7	1.2	293	0.591
	0 - 15	196	2.7	529	0.400	230	0.9	207	0.744
	15 - 30	667	0.53	354	0.315	280	2.5	700	0.794
	30 - 45	110	5.05	556	0.509	250	1.2	300	0.692
Mean		324	2.74	479.7	0.408	253	1.5	402	0.74
Umuokanne	0 - 15	105	5.01	526	0.461	240	1.3	312	0.910
	15 - 30	417	0.60	250	0.080	250	1.5	375	0.868
	30 - 45	435	1.28	557	0.575	270	2.6	702	0.560
Mean		319	2.30	444	0.372	253	1.8	463	0.779

b= Maximum adsorption, k=Affinity coefficient, MBC=Maximum buffering capacity, R²=Correlation coefficient, a=NH₄⁺-sorption capacity, n= NH₄⁺-sorption energy

(324mg/kg) and it increased with soil depth. Generally, the energy coefficient (k and n) increased with the adsorption maximum. This agrees with the earlier report by Wissen and Mongi (2015), who reported direct relationship between ammonium sorption maximum and energy coefficient. The correlation between ammonium sorption maximum, energy coefficients and some soil properties (Table 4) indicated that adsorption maxima and energy coefficients correlated negatively and significantly with clay content and pH. The clay fraction of the soil is dom-

inated by varying types of minerals especially 2:1 clay minerals which have high affinity for cations such as ammonium (Brady and Weil, 2008). Thus, the higher the clay constituent of a given soil, the more it's potential to adsorb nutrient elements such as ammonium. This agrees with the report of Wang *et al.*, (2000), who observed that ammonium sorption increases with clay content.

The energy coefficients (k and n), describes the ability and the tenacity to which ammonium ions are adsorbed

Table 4. Correlation between Langmuir and Freundlich constants and some soil properties

	Sand	Clay	pH	Organic Matter
b	-0.89	0.844	-1.000**	-0.567
k	0.944	0.728	-0.976	-0.401
a	-0.984**	0.827**	-0.998**	-0.540
n	-0.764	0.435	-0.834	-0.047

to clay mineral surfaces of the soils (Brady and Weil, 2008). The more the energy coefficients, the more the tenacity of ammonium adsorption and the less likely ammonium will be available in soil solution (Wang *et al.*, 2000). For Langmuir model, the k values increased with depth. This agrees with the earlier report by Wissen and Mongi (2015), who reported direct relationship between ammonium sorption maximum and energy coefficient. The higher the depth of the soil, the more the occurrence of mineral reactive surfaces thus the greater the

affinity coefficient for ammonium adsorption. FUTO had the greatest affinity for ammonium. The implication may be advantageous owing to the fact; leaching ammonium nitrogen may be reduced due to sorption. However, to ensure adequate ammonium nitrogen utilization by plants in this location, large and continuous ammonium fertilizer should be applied to compensate for the adsorbed fraction. Ihiagwa and Umuokanne that had lesser energy coefficients may be presented with more leaching of nitrogen. Sand and organic matter correlated neg-

atively with the affinity coefficients, while clay correlated positively with the energy coefficients (Table 4).

Considering the values of the R^2 for both Langmuir and Freundlich equations, it can be deduced that the Freundlich equation described the sorption parameters than the Langmuir equation. Comparing the two equations, it can be established that the Freundlich model showed a better fit to the data than the Langmuir model. The Freundlich isotherm provided the information about the surface heterogeneity and the exponential distribution of the active sites, their energies, and predicted the saturation of the surface of the sorbent by the sorbate. This result however contradicted the report of Wissen and Mongi (2015), who posited that Langmuir better fitted the sorption parameters of ammonium. The Langmuir poorly fitted the ammonium sorption data because of the heterogeneity the energy of the absorbent surfaces.

4.0 Conclusion

This study evaluated the ammonium sorption characteristics of soils from three locations (FUTO, Ihiagwa and Umuokanne) in Owerri west at three sampling depths (0-15cm, 15-30cm and 30-45cm). There was no significant variability on the soil physical and chemical properties of the soils with respect to location. However, soils from Ihiagwa consistently showed better fertility status compared with the other soils. The role of depth as it affects soil properties was significant. While soil pH did not follow a particular occurring trend, other nutrients generally decreased with depth.

Results of ammonium sorption study showed variability of ammonium sorption characteristics of the soils from different locations. The maximum sorption capacity of ammonium was obtained from soil of Ihiagwa (Mean = 324mg/kg), while the highest value of energy coefficient (5.03mg/kg) was obtained in soils of FUTO.

Generally, both the adsorption maxima and energy coefficients increased with soil depth. Results also indicated that soil properties, including pH, organic matter, amounts of sand and clay, significantly influenced the sorption of NH_4^+ by soils. Langmuir and Freundlich equations were applied to test the equilibrium data. Freundlich equation better described the sorption data, thus indicating the heterogeneity in the ammonium sorption characteristics of the different soil sampled.

References

- Azu, D.E.O., Osodeke, V.E. and Nwanja O.U. 2017. Effect of algae on phosphorus sorption characteristics of an ultisol of Southeastern Nigeria. *Direct Research Journal of Agriculture and Food Science*. (2):88-9.
- Azu, D.E.O., Osodeke, V.E., Ukpong, I.M. and Osi, A.F. 2018. Chemistry and mineralogy of soils derived from different parent materials in Southeastern Nigeria. *Int. J. of Plant and Soil Science*, 25 (25):1-16.
- Brady, N.C. and Weil, R.R. 2008. Element of the nature and properties of soils. Abridged. of: The nature and properties of soil, 13th ed.
- Carter, M. R. and Gregorich, E. G. 2008. Soil Sampling and Methods of Analysis 2nd (ed). Can Soc. Soil Sci. 1224p.
- Fox, R. L. and Kamprath, E.J. 1970. Phosphate sorption isotherms for evaluating the phosphate requirement of soils. *Soil Sci. Soc. Am. Proceeding*, 34:902-907.
- Gee, G.W. and D. O. (2002). Particle size distribution. In Dane, J.H. and G.C. Topp (eds).
- Methods of soil analysis. Part 4. Physical and mineralogical methods. Soil Science Society of America book series. No. 5 ASAS and SSA. Madison, Wisconsin. p 255-293. *Geoderma* 110: 191-204.
- NIMET (Nigerian Meteorological Agency), 2008. Climate Weather and Water Information, for Sustainable Development and Safety
- Onwuka, M.I., Osodeke, V.E. and Okolo, N.A. 2007. Amelioration of soil acidity using cocoa husk ash for maize production in Umudike area of southeastern Nigeria. *Tropical and Subtropical Agroecosystem* 7:41-45
- Onyegbule, O., Azu, Donatus, E.O., Ike, S. and Akagha, U. 2018. Comparing organic matter content under four different land uses in Uloandugba, Imo State, Nigeria. *Journal of Applied Life Science, International*, 22(2):1-6.
- Osodeke, V. E. 1996. Effects of rainfall and land-use on the management of the highly degraded soils of southeastern Nigeria. *Proceedings of the 5th Annual conference of the Nigerian Society of Biological conservation*, Umudike.
- Osodeke, V. E. and Uba, A. F. 2005. Determination of phosphorus fraction in selected soils of southeastern Nigeria. *International Journal of Natural and Applied Sciences (IJNAS)*, 1(1):10-14.
- Osodeke, V.E. 2005. Effect of phosphorus on nodulation and yield of soybean (*Glycine max*) in soils of the forest zone of southeastern Nigeria. *J. Appl. Chem. Agric. Res.* 7:170-177.
- Pansu, M. and Gautheyrou, J. 2006. Handbook of Soil Analysis: Mineralogical Organic and Inorganic Methods. Springer, p995.
- Ranjbar F, Jalali M. 2013. Measuring and modeling ammonium adsorption by calcareous soils. *Environ. Monit. Assess.* 185: 3191-3199. doi.org/10.1007/s10661-012-2782-y
- Simmons, E. N., Jones, J. B., Mills, A. H., Smittle, A. A. and Hussey, C. G. 1994. Comparison of analytical methods for nitrogen analysis in plant tissues. *Commun Soil Science*, 24:1609-1616
- Sun, Zhao, Allen 2005. Enhanced removal of organic matter and ammoniacal-nitrogen in a column experiment of tidal flow constructed wetland system. *J. Biotechnol.* 115: 189-197. doi.org/10.1016/j.jbiotec.2004.08.009.
- Sutton M.A., Van-Grinsven H., Grizzetti B., Howard C.M., Erisman J.W., Billein G., Bleeker A., Grennfelt P. (2011a): The European Nitrogen Assessment Cambridge university press Cambridge.
- Tyrrel, S.F., Harrison, P.B.L., Harrison, K.S. 2002. Removal of ammoniacal nitrogen from landfill leachate by irrigation onto vegetated treatment planes. *Water Res.* 36: 291-299. doi.org/10.1016/

S0043-1354(01)00217-2.

Udo. E.J. T.O. Ibia, J.O.O. gunwale, A. O. Anoan-
dI.Esu.2009. *Manual of Soil, Plant and Water Analysis*.
Sibon Books Ltd. Lagos.

Onweremadu, E.U. and Ufot, U.O. 2011. Soil nitrogen
forms distribution in isohyperthermic kandiodiults of
central southeastern Nigeria. *Nigeria journal of Agri-
culture, food and environment*, **7**(2), 52-56

Wang MY, Glass A, Shaft JE and Kochians LV. 2000.
Ammonium uptake by rice roots (III. Electro physi-
ology). *Plant physiology* 104:899-906

Wissem, H., Jean, A. M., David, P., Noura, Z. and Mongi,
S. 2015. Phosphorus isotherm sorption in semi-arid

soil. *International Journal of Engineering and Tech-
nology*, 2(3):1995-2005.

Zhang, S., Zhang, X., Huffman, T., Liu, X. and Yang, J.
2011. Influence of topography and land management
on soil nutrients variability in Northeast China. *Nu-
trient Cycling Agroecosystem*, 89, 427-43.

Zhu, W. L., Cui, L. H., Ouyang, Y., Long, C. F., & Tang,
X. D. 2011. Kinetic adsorption of ammonium nitro-
gen by substrate materials for constructed wet-lands.
Pedosphere. 21: 454-463. doi.org/10.1016/S1002-
0160(11)60147-1.