




Nigerian Journal of Soil Science

Journal homepage: www.soilsjournalnigeria.com

Dynamics in physico-chemical properties of soils under oil palm plantations of different ages

Olufemi Adewale Osinuga

Department of Soil Science and Land Management, Federal University of Agriculture, Abeokuta, Nigeria

ARTICLE INFO

Article history:

Received June 20, 2021

Received in revised form July 13, 2021

Accepted August 20, 2021

Available online September 12, 2021

Keywords:

Climate change

Forests

Nutrient dynamics

Oil palm plantations

Tropical soils

ABSTRACT

This research investigated the changes in the physicochemical properties of soils under oil palm plantations of different ages. Soil surface (0-20 cm) and subsurface (20-40 cm) samples have been obtained from various ages of oil palm plantations (0-5, 5-10, 10-15 and, 15-20 years). Two distinct samples were taken on the same farm, under alleys and heaped pruned fronds. Soil samples used as a standard (control) were collected from adjacent forest land. Analyses of particle size showed that the soils were sandy loam to sandy clay loam texture soils. Bulk density was low to moderate ($0.93 - 1.25 \text{ g/cm}^3$), and varied with age and depth. The soils were moderate to neutral pH ($5.38 - 6.81$), low to moderate organic carbon ($11.6 - 21.7 \text{ g/kg}$) and total nitrogen ($1.18 - 2.03 \text{ g/kg}$), and relatively low available phosphorus ($4.43 - 6.21 \text{ mg/kg}$) contents. Based on the standard ratings, cation exchangeable capacity content was low ($4.93 - 6.15 \text{ cmol/kg}$), while high percent base saturation ($85.3 - 93.2\%$) was observed. The soil properties determined showed that soil nutrients under alleys fluctuated with age whiles that under heaped fronds increased. Research findings have shown that the soil properties of different ages of oil palm plantations vary and should be handled differently based on their characteristics. Accumulation of organic residue on the floor of the plantations should be encouraged as this will help increase organic matter levels.

Corresponding Author's E-mail Address:

osinugaoa@funaab.edu.ng

ISSN– Online **2736-1411**

Print **2736-142X**

© Publishing Realtime. All rights reserved.

1.0 Introduction

Oil palm (*Elaeis guineensis*) is one of the plantation crops grown in some tropical countries of the world especially Nigeria in addition to cocoa, kola, teak, gmelina, rubber, and cashew a result of transformations in agriculture. It is cultivated mainly in the southern, eastern, and western parts of the country. Recently, planted areas have increased at a rapid pace, and the increase is significantly evident. Oil palm was adopted and cultivated among the high-value perennial tree crops by individual small-scale growers, private agencies, and government agencies (state and federal). Global oil palm production is approximately 272,055,131 tonnes from 18,917,400 hectares, with Asia (84.1%) taking the lead, followed by Africa (9%) (FAO, 2020). It is a major source of oils and fats for human food, livestock feed, and the manufacturing of several other domestic products, such as cosmetics, soap, and detergents (Reiger, 2006).

The increase in oil palm demand has resulted in the conversion of natural vegetation into oil palm plantations. The conversion of some rainforests and the use of abandoned logged-off forests into oil palm plantations is expected to have contributed significantly to nutrient losses from both soil and vegetation (Brahene *et al.*, 2016; Rozieta *et al.*, 2016). Deforestation and loss of habitat of critically endangered species (Clay, 2004), a decrease in soil productivity, an increase in soil erosion, and soil biodiversity loss (Comte *et al.*, 2012; Savilaakso *et al.*, 2014), and a substantial increase in greenhouse gas emissions are the detrimental effects of oil palm on the climate (Bates *et al.*, 2008; Hassan *et al.*, 2011). Rising tillage intensity and the conversion of the natural environment to agricultural land have contributed to a decline in soil organic matter levels due to decreased organic carbon inputs and decreased physical conservation of soil organic carbon contents (Chibsa and Ta', 2009).

As a consequence of the combined effects of physical,

chemical, and/or biological processes operating at various intensities and on different scales, the classification of soils depends on the degree of spatial heterogeneity (Priyabrata *et al.*, 2008). Research has shown that crop age also leads to soil variability because nutrients are extracted from the soil during harvest for grain, fibre, wood, and crop residues as the crops grow older (Basiron, and Weng, 2004; Aweto and Enaruvbe, 2010). If replenishment with inorganic and organic fertilizers is insufficient, nutrient removal can result in a decrease in soil fertility (Okon *et al.*, 2017). There seems to be little or no knowledge about the impact of oil palm plantations of different ages on soil physico-chemical properties in the research region to date on various studies on oil palm plantations in Nigeria. Therefore, it is of great importance to assess the quality of soil nutrients in existing oil palm plantations of various maturity ages and to analyze the variability (differences) in the soil properties of these oil palm plantations.

2.0 Materials and methods

2.1 Site Description

Oil palm plantations owned by the Federal University of Agriculture, Abeokuta (FUNAAB) and Ogun State Ministry of Agriculture, were selected with a reference (control) soil adjacent to the plantations (Figure 1). FUNAAB is located next to Ogun-Oshun River Basin Development Authority (OORBDA), along Alabata road, while the ministry is along Abeokuta-Ibadan expressway in Odeda Local Government Area (LGA) of Ogun State. Odeda lies in a humid tropical lowland zone with two separate seasons (wet and dry). The wet season runs between March and October, and the dry season runs between November and February. The annual rainfall is between 1000-1500 mm, the annual temperature is between 26-32°C, and the relative humidity varies between 70-88%. The LGA superimposes the Basement Complex's pre-Cambrian metamorphic rocks with bed-rock consisting mainly of granitic gneisses, horn-blended gneisses, bounded biotite, quartzite, and quartz schists.

2.2 Field Work and Sample Collection

The oil palm plantations ranged from young ones aged around five years to farms as old as twenty years at various maturity ages. Marking an area of 30 m by 30 m was followed by sampling. The sampling of farms commenced by establishing four age-based clusters of oil palm plantations into which numerous farms were clustered. A total of twelve parcels, with three parcels for each age group, were chosen as replicates for sampling. Soil samples were obtained at depths of 0-20 cm and 20-40 cm. There were 0-5, 5-10, 10-15, and 15-20 years for the different plantation age groups considered. The sampled plots comprised both alleys within the palm rows and pruned and heaped palm fronds. For bulk density analysis using core samplers, undisturbed soil samples were collected within the alleys and under heaped trees. Under the pruning, especially under old heaps, sampling was conducted with care because the top layer had to be separated from the decomposed material sitting just above it. Soils from the sampled spots of various depths from each plantation were put together to obtain a composite sample. A sub-sample was taken, air dried, crushed, and sieved through a 2 mm sieve for routine laboratory testing, and processed.

2.3 Laboratory Analysis

Using the hydrometer method (Gee and Or, 2002), particle size composition was performed, bulk density was deter-

mined using the core sample method. The pH was calculated electrometrically using the glass electrode pH meter in soil-water suspension. The soil organic carbon was measured using the digestion method of wet oxidation (Walkley and Black, 1934) and total nitrogen by the digestion method of macro-Kjeldahl (Bremmer and Mulvaney, 1982). The Bray-1 extractant was used to extract available phosphorus (Bray and Kurtz, 1945), while the P in the extract was determined by the vanado-molybdate blue method (Murphy and Riley, 1962). With 1.0 M KCl and titrated with NaOH, exchangeable acidity (H^+ and Al^{3+}) was extracted, and with 1.0 M NH_4OAc at pH 7, exchangeable bases were extracted. The atomic absorption spectrophotometer was used to determine Ca^{2+} and Mg^{2+} , while the flame photometer read K^+ and Na^+ . Cation Exchangeable Capability (CEC) was obtained by the summation of exchangeable bases and total acidity (Chapman, 1965). The base saturation was obtained as the ratio of exchangeable bases to CEC.

2.4 Statistical Analysis

Data were analyzed using descriptive statistics to show the relationship between the variables in the plantations. The mean was used to derive the average distribution of the variables, the standard deviation shows how the variables deviate from the mean.

3.0 Results

The physical and chemical properties of the soils are depicted in Tables 1 and 2. The particle size fractions varied significantly ($p < 0.05$) with the age of plantations, and also varied with depths. The sand fractions which ranged between 698-778 g/kg (Table 3) were higher in the oil palm plantations, while silt (50-80 g/kg) and clay (172-232 g/kg) fractions were higher in the forest (uncultivated) soils. Analyses of the particle size revealed that the soil texture was sandy loam and sandy clay loam. Bulk density (BD) values ranged between 0.93-1.25 g/cm³ and increase with depth. Under both alleys and heaps, the rise in bulk density with depth was noticed, and this could be due to the increasing clay content with depth. In the 0-20 cm layer, soil bulk density levels are relatively lower than those in the 20-40 cm layer.

The soil pH values ranged from moderate to slightly acidic except for the reference soil which was neutral. The pH was higher in the forest surface layer with a value of 7.13 (Table 1) and lowest at the oil palm alley 15-20 years with a value of 5.74 (Table 2). Between the 0-20 and 20-40 cm layers, the results varied significantly. Inside the oil palm alleys, the pH of the soil appears to be lower than that under heaped fronds and fluctuates as the years of planting increase. Generally, available P did show a significant difference across all the oil palm plantations and soil depths at $p < 0.05$. The average available P in the topsoil was higher than in the subsoil. The highest value of the uncultivated topsoil was 6.21 mg/kg, followed by 6.10 mg/kg for the oil palm heaps 15-20 years (Table 1), with the lowest values being 4.43 mg/kg in the oil palm alley 5-10 years (Table 2).

The soil organic carbon (OC) content was moderate, ranging from 12.4 g/kg to 21.7 g/kg for 0-20 cm depth (Table 1) while at 20-40 cm depth, it ranged from 11.6 g/kg to 20.4 g/kg (Table 2). The OC content of the reference (uncultivated) site was significantly ($p < 0.05$) higher than the cultivated sites. Among the oil palm plantations, the 15

-20 years under heaps have the highest OC content (17.8 g/kg), while the 1-5 years have the lowest OC content (11.6 g/kg). The reference (uncultivated) site's OC content was significantly ($p < 0.05$) higher than the sites under cultivation. The OC remained constant for some years after the removal of the current oil palm forest vegetation cover, but after 10 years it was observed that OC accumulation occurred in both alleys and under heaped fronds, especially at a depth of 0-20 cm, aside from that the values observed under heaped fronds were slightly higher. The soil's total nitrogen (TN) was moderate and ranged from 1.18-2.03 g/kg. The oil palm plantation of 0-5 years and 5-10 years under alley has the same value of 1.18 g/kg at the subsoil. The overall distribution of nitrogen content followed a similar trend to the OC distribution and differed significantly with the age of the plantation.

Exchangeable acidity (EA) was low and fluctuated with the years of oil palm cultivation. No substantial variation was found between the overall EA and the age of the plantations and the depth of the soil at $p > 0.05$. In the uncultivated soil at both depths, the exchangeable bases were higher than in the different age groups of the oil palm soils, both in the alleys and under the palm fronds (Tables 1 and 2). According to the Federal Fertilizer Department's ranking (2012), exchangeable K is low to moderate in both reference soils and oil palm soils. With respect to the soil depth, the mean average K also showed no noticeable difference but declined with the depth of the soil. The soil's exchangeable Na is low and the amount seems too small to raise some concerns about any potential physical effect of the soil. Compared to the remaining bases, exchangeable Ca values were greater, followed by exchangeable Mg. In general, exchangeable Mg values initially appeared to increase with time in both alleys and under heaped fronds, but declined gradually with further planting age, while the other bases (Ca, Na and K) fluctuated with the age of the plantations.

Cation Exchange Capacity (CEC) did show significant variation across the plantations at $p > 0.05$. There was also significant variation in CEC between the depths (Table 1 and 2). However, at a depth of 0-20 cm, the reference soils had the highest CEC value of 6.15 cmol/kg followed by the 0-5 years plantation the under alleys with 5.50 cmol/kg while the 5-10 years plantation under heaps had the lowest value of 5.01 cmol/kg. The soil CEC was low (< 16 cmol/kg) and decreased with depth. There was a general increase in the CEC in both layers after 10-15 years of the alleys and heaped palm fronds to the soil. The percent base saturation (BS %) of the soils was greater than 50%, and the trend is somewhat inconsistent with the plantation age. The values ranged between 85.3% and 93.2% (Table 3).

The calculated correlation coefficients used for assessing the relationships among soil variables are presented in Table 4. The sand is the dominant fraction correlated negatively with all the soil parameters except for H and Al, whereas an inverse correlation (positive) was observed with the clay and silt contents. Soil pH is correlated positively and significantly with OC ($r = 0.919^{**}$), TN ($r = 0.873^{**}$) and Av. P ($r = 0.638^{**}$), and negatively with H ($r = -0.442^{**}$) and Al ($r = -0.498^{**}$), indicating that pH as a master variable regulates the availability of basic cations in the soil. Soil OC was also positively related to TN ($r = 0.974^{**}$) and Av. P ($r = 0.750^{*}$). TN show positive relationship with Av. P ($r = 0.742^{**}$) and negative with both H and Al ($r = -0.460^{*}$ and $r = -0.485^{**}$). The Av. P related positively with all the basic cations (Na, K, Ca, and Mg), but negatively correlated with basic anions (H and Al). The CEC correlated perfectly with BD ($r = 0.355^{**}$), pH ($r = 0.442^{**}$), OC ($r = 0.370^{**}$), TN ($r = 0.333^{**}$), Av. P

($r = 0.180^{**}$) and the basic cations, while it showed negative correlation with H ($r = -0.523^{**}$) and Al ($r = -0.069$).

4.0 Discussion

The physical and chemical properties of the soil in different land use may be attributed to differences in the study area's management practices. Soil texture is an inherent property; spatial variations between soils with less variation associated with cultivation could be due to the textural differences found (Brahene *et al.*, 2016). In the 0-20 cm layer, soil bulk density values were comparatively lower when compared to those of the 20-40 cm layer where the content of organic matter was very low. This suggests that OM has contributed significantly to enhance the soil's physical properties, thereby contributing to the soil's structural stability (Germer and Sauerborn, 2008). The BD values obtained cannot hinder root development and penetration.

The low pH of the soils can be attributed to the nature of the parent materials and the high precipitation that causes the basic cations in the soil to leach intensively (Tweneboah, 2000; Owusu-Bennoah *et al.*, 2000; Oyegoke *et al.*, 2017). Low pH values connote the presence of a positively charged colloidal surface capable of attracting negatively charged ions. However, the values for oil palm production were acceptable because they were below 7.5 and did not favour oil palm production above that value (Okon *et al.*, 2017). Available soil phosphorus level was low, as the values were between 3-7 mg/kg (Federal Fertilizer Department, 2012). This means that the amount of phosphorous is according to the pH status of the oil palm plantations. In this land-use system, the higher available P content within the forest may be correlated with increased microbial activity.

The high OC content of soils under heaped fronds is related to the quantity, position, consistency, and temperature and humidity actions on the pruned fronds compared to those under alleys (Kirschbaum, 2000; Comte *et al.*, 2012). As a result of the rapid decomposition of palm fronds, there could be a rise in OC and soil nutrients in planting for less than 10 years, but as the planting age moved to 25 years, the decomposition rate slowed down with decomposed material being covered by overlying palm fronds (Okon *et al.*, 2017). Thus, with the age of the plantation, it is possible to find heaps of different heights (Brahene *et al.*, 2016). Total nitrogen (TN) varies with the quantity of organic matter present in soils, so it has risen without exception, along with improvements in the related status of organic matter. Relatively, TN amount is also determined by organic carbon, which in turn results from plant and root biomass, as well as residues returned to the soil system. The fundamental cause of N deficiency in tropical soils is extreme leaching and erosion due to high tropical precipitation (Aweto and Enaruvbe, 2010; Osinuga and Oyegoke, 2019).

The low level of basic cations in the plantations results from the effect over time of the continuous uptake of nutrients by the plants. The result shows that the intensity of weathering, cultivation, and use of inorganic acid-forming fertilizers affects the distribution of the cations in the soil system and improves their depletion (Owusu-Bennoah *et al.*, 2000). The low values observed for these basic cations may also be due to the low content of organic matter and the presence of low activity clays of the area (Oyegoke, 2011). Subsequently, the obtained CEC is a function of the pH and SOM in the soil and, with the age of the oil palms, the value remained reason-

ably constant in the soil. This is an indication that the soils remain low in CEC at their natural pH levels, demonstrating the soils' low nutrient retention ability.

Sand negative interaction with clay connotes that an increase in one is a decrease in the other. The relationship bulk density enjoys with silt and clay portray particle size's influence on this soil property. Soil pH is interrelated with organic carbon and total nitrogen, indicating that pH as a master variable regulates the decomposition of organic materials and availability of nitrogen in the soil. Organic matter content is a sink and a source for many plant nutrients exhibits significant

relationship with total nitrogen and available phosphorus. Similarly, total N correlated significantly with available P. The close to unity, that is, the perfect relationship exhibited by total N with organic C content implies that almost 94% of total N in the soil are derived from microbial decomposition and mineralization of organic residues. Mg being the second dominant cation in the absorption complex; is positively and highly significantly related to Na, K, Ca and CEC. Available P interrelated weakly with the basic anions (H and Al) owing to the fact that phosphorus availability had a significant effect on basic anions in the soil. Nearly all the soil variables interrelated very well the CEC. This might be at-

Table 1: Physical and Chemical Properties of Soils under Oil Palm Alley and Heaps (0-20 cm)

Plantation	Sand	Silt	Clay	BD	pH	OC	TN	Av. P	H ⁺	Al ³⁺	Ca	Mg	Na	K	CEC	BS
Age (Years)	g/kg.....			g/ cm ³	(H ₂ O)	g/kg	g/kg	mg/kg	 cmol/kg							%
Alley and Reference Soil																
0-5	758	60	182	1.06	6.07	12.4	1.23	5.67	0.03	0.54	2.69	1.51	0.44	0.29	5.50	89.6
5-10	778	50	172	1.02	5.91	12.6	1.24	4.89	0.04	0.42	2.57	1.58	0.39	0.26	5.26	91.3
10-15	748	70	182	1.13	5.98	15.2	1.41	5.64	0.04	0.46	2.43	1.78	0.33	0.28	5.37	90.7
15-20	728	70	202	0.98	5.79	15.9	1.42	5.80	0.06	0.39	2.58	1.53	0.41	0.32	5.29	91.5
Uncultivated	698	80	222	1.23	7.13	21.7	2.03	6.21	0.02	0.40	2.86	2.02	0.51	0.34	6.15	93.2
Heaps and Reference Soil																
0-5	748	70	182	1.04	6.14	12.5	1.24	5.69	0.04	0.48	2.52	1.38	0.45	0.31	5.18	90.0
5-10	758	60	182	0.93	6.03	13.1	1.26	5.73	0.05	0.50	2.50	1.23	0.41	0.27	5.01	89.1
10-15	748	70	182	1.11	6.08	16.5	1.79	6.02	0.03	0.57	2.48	1.42	0.34	0.30	5.14	88.3
15-20	718	70	212	1.09	5.88	17.8	1.84	6.10	0.04	0.52	2.45	1.50	0.31	0.28	5.11	86.9
Uncultivated	698	80	222	1.23	7.13	21.7	2.03	6.21	0.02	0.40	2.86	2.02	0.51	0.34	6.15	93.2

BD = Bulk density; TN = Total Nitrogen; OC = Organic carbon; Av. P = Available Phosphorus; CEC = Cation exchange capacity; BS = Base saturation.

Table 2: Physical and Chemical Properties of Soils under Oil Palm Alley and Heaps (20-40 cm)

Plantation	Sand	Silt	Clay	BD	pH	OC	TN	Av. P	H ⁺	Al ³⁺	Ca	Mg	Na	K	CEC	BS
Age (Years)	g/kg.....			g/ cm ³	(H ₂ O)	g/kg	g/kg	mg/kg	 cmol/kg							%
Alley and Reference (Uncultivated Forest) Soil																
0-5	748	70	182	1.08	6.03	11.6	1.18	5.26	0.06	0.63	2.43	1.37	0.35	0.26	5.10	86.5
5-10	768	60	172	1.03	5.86	11.7	1.18	4.43	0.05	0.72	2.48	1.56	0.25	0.19	5.25	85.3
10-15	728	80	192	1.16	5.92	12.5	1.35	5.14	0.02	0.55	2.37	1.60	0.31	0.24	5.09	89.2
15-20	718	70	212	1.12	5.74	13.3	1.37	5.24	0.07	0.61	2.38	1.62	0.28	0.21	5.17	86.9
Uncultivated	688	80	232	1.25	7.01	20.4	1.92	5.82	0.03	0.49	2.75	1.79	0.40	0.29	5.75	91.0
Heaps and Reference (Uncultivated Forest) Soil																
0-5	748	70	182	1.08	6.11	12.1	1.20	5.27	0.03	0.58	2.44	1.34	0.33	0.27	4.99	87.7
5-10	758	60	182	1.04	6.02	12.7	1.21	5.48	0.04	0.65	2.51	1.43	0.26	0.22	5.11	86.5
10-15	728	70	192	1.15	6.06	15.4	1.49	5.73	0.03	0.61	2.50	1.52	0.28	0.21	5.15	87.6
15-20	718	70	212	1.11	5.85	16.3	1.58	5.77	0.06	0.42	2.39	1.60	0.19	0.27	4.93	90.3
Uncultivated	688	80	232	1.25	7.07	20.4	1.92	5.82	0.03	0.49	2.75	1.79	0.40	0.29	5.75	91.0

BD = Bulk density; TN = Total Nitrogen; OC = Organic carbon; Av. P = Available Phosphorus; CEC = Cation exchange capacity; BS = Base saturation.

Table 3: Descriptive Statistics of Soil Samples from the Oil Palm Plantations

Variable	Descriptive Statistics									
	N	Range	Minimum	Maximum	Means	Std. Error	Std. Deviation	Variance	Skewness	Kurtosis
Sand (g/kg)	20	90	688	778	733.50	5.959	26.651	710.263	-.312	-.881
Silt (g/kg)	20	30	50	80	69.00	1.762	7.881	62.105	-.531	.490
Clay (g/kg)	20	60	172	232	196.50	4.441	19.861	394.474	.597	-1.092
BD	20	0.32	0.93	1.25	1.1045	0.01989	0.08894	0.008	.123	-0.427
pH	20	1.27	5.86	7.13	6.2770	0.09825	0.43939	0.193	1.349	0.225
OC (g/kg)	20	10.10	11.60	21.70	15.2900	0.77269	3.45557	11.941	0.795	-0.720
TN (g/kg)	20	0.85	1.18	2.03	1.4945	0.06918	0.30939	0.096	0.687	-1.153
Av. P (mg/kg)	20	1.78	4.43	6.21	5.5960	0.10049	0.44941	0.202	-0.929	1.012
H ⁺ (cmol/kg)	20	0.05	0.02	0.07	0.0395	0.00328	0.01468	0.000	0.539	-0.581
Al ³⁺ (cmol/kg)	20	0.33	0.39	0.72	0.5215	0.02088	0.09337	0.009	0.332	-0.621
Ca ²⁺ (cmol/kg)	20	0.49	2.37	2.86	2.5470	0.03434	0.15356	0.024	0.981	-0.158
Mg ²⁺ (cmol/kg)	20	0.79	1.23	2.02	1.5795	0.04724	0.21125	.045	0.717	0.226
Na ⁺ (cmol/kg)	20	0.32	0.19	0.51	0.3575	0.01917	0.08571	.007	0.080	-0.435
K ⁺ (cmol/kg)	20	0.15	0.19	0.34	0.2720	0.00936	0.04188	.002	-0.295	-0.386
CEC (cmol/kg)	20	3.32	2.83	6.15	5.2175	0.14761	0.66013	.436	-2.376	9.462
BS (%)	20	7.90	85.30	93.20	89.2900	0.50855	2.27432	5.173	0.043	-0.879
Valid N (listwise)	20									

BD = Bulk density; TN = Total Nitrogen; OC = Organic carbon; Av. P = Available Phosphorus; CEC = Cation exchange capacity; BS = Base saturation.

Table 4: Pearson Correlation Coefficients Among the Soil Properties in the Oil Palm Plantations

	Sand	Silt	Clay	BD	pH	OC	TN	Av. P	H	Al	Na	K	Ca	Mg	CEC
Sand	1.000														
Silt	0.900*	1.000													
Clay	0.974*	0.804*	1.000												
BD	0.813*	0.803*	0.742*	1.000											
pH	0.861*	0.783*	0.879*	0.767*	1.000										
OC	0.850*	0.767*	0.857*	0.742*	0.919*	1.000									
TN	0.837*	0.762*	0.833*	0.764*	0.873*	0.974*	1.000								
Av. P	0.612*	0.618*	0.594*	0.400	0.638*	0.750*	0.742*	1.000							
H	0.263*	-0.323	-0.172	0.554*	0.442*	-0.430	0.460*	-0.344	1.000						
Al	0.398*	-0.305	0.447*	-0.197	0.498*	0.571*	0.485*	0.576*	0.181	1.000					
Na	0.252*	0.308	0.254*	0.202	0.547*	0.435	0.367*	0.466	0.457*	0.571*	1.000				
K	0.411*	0.501*	0.413	0.267	0.613*	0.618	0.574*	0.741*	-0.366	0.796*	0.787*	1.000			
Ca	0.505*	0.432	0.545*	0.504*	0.819*	0.721*	0.631*	0.494	-0.542	-0.483	0.788*	0.634*	1.000		
Mg	0.683*	0.597*	0.684*	0.789*	0.748*	0.775*	0.719*	0.341*	0.411*	0.486*	0.378	0.395*	0.669*	1.000	
CEC	0.230*	0.288*	0.202*	0.355*	0.442*	0.370*	0.333*	0.180*	0.523*	-0.069	0.721*	0.327*	0.675*	0.453	1.000

* Significant at the 0.05 level; ** Significant at the 0.01 level.




Figure 1: Location Map of Study Sites

5.0 Conclusions

The study concludes that the distribution of particle size of the soils studied did not vary significantly because they were formed from the same parent material (basement complex rocks). Nutrient mining in plantations is possible as continuous cultivation has been observed to minimize major and minor plant nutrients. Heaping palm fronds were found to yield some advantages over time in terms of carbon content, but could not provide enough nutrients to substitute what the crop used. The rate of decomposition of organic materials increased with high temperatures, and the release of nutrients was faster than plants could quickly absorb and use, and was often subject to losses associated with erosive precipitation. As pruned fronds and cut plants continue to boost the structure of the soils to the point that soil bulk density values are relatively low, the impact of organic matter addition to soils was seen to be beneficial. The study recommends that the accumulation of organic residue on the floor of plantations should be championed as it will help sustain increasing levels of organic matter. It is important to promote more research on how to efficiently use pruned fronds for compost to be spread around the oil palm trees.

Acknowledgements

The author wishes to appreciate the Directorate of University Farms, Federal University of Agriculture Abeokuta and Ogun State Ministry of Agriculture, who allowed me to use their oil palm farms and necessary support rendered in the course of the fieldwork.

References

- Aweto, A.O. and Enaruvbe, G.O. (2010). Catenary variation of soil properties under oil palm plantation in South-Western Nigeria. *Ethiopian J. Envntal. Stud. & Mgt*, 3(1):1-10.
- Basiron, Y. and Weng, C.K. (2004). The oil palm and its sustainability. *J. Oil Palm Res.* 16(1):1-10.
- Bates, B.C., Kundzewicz, Z.W., Wu, S. and Palutikof, J.P. (2008). Climate change and water. Technical paper of the Intergovernmental Panel on climate change, IPCC Secretariat, Geneva, 210pp.
- Brahene, S.B., Owusu-Bennoah, E and Abekoe, M.K. (2016). Physico-chemical properties of soils under oil palm plantations of different ages. *Nature & Faune* 30(1): 54-58.
- Bray, R.H. and Kurtz, L.T. (1945). Determination of total organic and available forms of phosphorus in soils. *Soil Sci* 59: 222-229.
- Bremner, J.M. and Mulvaney, C.S. (1982). Total Nitrogen. In: A. L. Page, R. H. Miller, and D. R. Keeney (Eds.), *Methods of Soil Analysis*. Madison, W.I. *Am Society Agron.* 2:49-158.
- Chapman, H.D. (1965). Cation exchange capacity. In: Black, C. A. (Ed.). *Methods of soil analysis*. American Society of Agronomy, Madison, pp 891-901.

- Chibsa, T. and Ta', A.A. (2009). Assessment of soil organic matter under four in Bale Highland, Southeast Ethiopia. Soil organic matter in four land use systems: forestland, grassland, fallow land and cultivated land. *World Appl. Sci. J.* 6(9): 1231-1246.
- Clay, J. (2004). Palm oil. In "World Agricultural and the Environment: A Commodity-by-Commodity Guide to Impacts and Practices" (Jason Clay, Ed.), Island Press, Washington, DC, USA, pp 203–235.
- Comte, I., Colin, F., Whalen, J.K., Grünberger, O., Caliman, J.P. and Gru, O. (2012). Agricultural practices in oil palm plantations and their impact on hydrological changes, nutrient fluxes and water quality in Indonesia: A review. *Adv. Agron.* 116: 71-122.
- Federal Fertilizer Department (FFD). (2012). Fertilizer use and management practices for crop production in Nigeria. 4th edn. Eds V. O. Chude; S. O. Olayiwola; A. O. Osho and C. K. Daudu, Federal Ministry of Agriculture and Rural Development, Abuja, Ni-geria. ISSN 115-554X.
- FAO. (2020). Food and agricultural data of the world. Production quantities of the oil palm fruit. <http://www.fao.org/faostat>. Accessed 13/10/2020.
- Gee, G.W. and Or, D. (2002). Particle size distribution. In: Dane, J.H., Topp, G.C. (Eds) Methods of soil analysis, part 4, physical methods. Soil Science Society of American Book Series No 5, ASA and SSSA, Madison, W.I., pp 255-293.
- Germer, J. and Sauerborn, J. (2008). Estimation of the impact of oil palm plantation establishment on greenhouse gas balance. *Envtal. Dev. Sust.* 10(6): 697–716.
- Hassan, M.N.A., Jaramillo, P. and Griffin, W.M. (2011). Life cycle GHG emissions from Malaysian oil palm bioenergy development: the impact on transportation sector's energy security. *Energy Policy* 39: 2615–2625.
- Kirschbaum, M.U.F (2000). Will changes in soil organic carbon act as a positive or negative feedback on global warming? *Biogeochem.* 48: 21–51.
- Okon, M.A., Nwachukwu, M.N. and Osujieke, D.N. (2017). Differences in physicochemical properties of soils under oil palm plantations of different ages in Ohaji/Egbema, Imo State. *Int'l J. Res. Agric. Forestry* 4(1): 1-5.
- Osinuga, O.A. and Oyegoke, C.O. (2019). Degradation assessment of wetlands under different uses: implications on soil quality and productivity. *Afri J. Agric. Res.* 17 (1): 10-17.
- Owusu-Bennoah, E., Awadzi, T.W., Boateng, E., Krog, L., Breuning-Madsen, H. and Borggaard, O.K. (2000). Soil properties of a toposequence in the moist semi-deciduous forest zone of Ghana. *West Afri. J. Appl. Eco.* 1: 1-10.
- Oyegoke, C.O. (2011). Clay mineralogy of major soils in geological transition zone; Isolu-Opeji, Ogun State, Nigeria. Unpublished PhD. Thesis University of Agriculture, Abeokuta, Nigeria.
- Oyegoke, C.O., Osinuga, O.A., Senjobi, B.A. and Martins, O. (2017). Effects of chemical fertilizers and herbicides on accumulation of some heavy metals in wetlands under different land use types in Odeda, Southwest Nigeria. *Moor J. Agric. Res.* 18: 1-12.
- Priyabrata, S., Chroppa, U.K. and Debashis, C. (2008). Spatial variability of soil properties and its application in predicting surface map of hydraulic parameters in an agricultural farm. *Current Sci.* 95(7): Sp 1.
- Reiger, M. (2006). Introduction to fruit crops. 6th edn. Harworth Press Inc., New York, pp 26-30.
- Rozieta, R., Sahibin, A. and Wan Mohd, R.I. (2016). Soil carbon sequestration in different ages of oil palm plantations. *Am-Eur. J. Agric. & Envthal. Sci.* 16(5): 1004-1007.
- Savilaakso, S., Garcia, C., Garcia-Ulloa, J., Ghazoul, J., Groom, M., Guariguata, M.R., Laumonier, Y., Nasi, R., Petrokofsky, G., Snaddon, J. and Zrust, M. (2014). Systematic review of effects on biodiversity from oil palm production. *Envtal. Evid.* 3: 4.
- Tweneboah, C.K. (2000). Modern Agriculture in the Tropics, with Special Reference to West Africa Cash Crops. Co-Wood Publishers, pp 121-147.
- Walkley, A. and Black, I.A. (1934). Determination of organic matter in soil. *Soil Sci.* 37: 549-556.