



# Nigerian Journal of Soil Science

Journal homepage: [www.soilsjournalnigeria.com](http://www.soilsjournalnigeria.com)

## Effects of Phosphorus on the Phenological Traits of Maize (*Zea Mays*) in Yola North Eastern Nigeria.

<sup>1</sup>Kwaghe, E.J. and <sup>2</sup>Gungula, D<sup>1</sup> Raw Materials Research and Development Council, Abuja, Nigeria<sup>2</sup> Department of Crop Production and Horticulture, Federal University of Technology, Yola

### ARTICLE INFO

#### Article history:

Received November 18, 2023

Received in revised form December 16, 2023

Accepted December 23, 2023

Available online January 21, 2024

#### Keywords:

Phosphorus  
Tasselling  
Anthesis  
Silking  
Maturity  
maize

### ABSTRACT

This study was conducted during the 2000 and 2001 cropping seasons at the University Research Farm, Federal University of Technology, Yola to evaluate the effects of phosphorus on the phenological traits of maize (New Kaduna White) hybrid variety. There were eight treatments consisting of four levels of N; 0, 60, 120 and 180kgNha<sup>-1</sup> and two levels of P, 0 and 60kgP<sup>-1</sup>, replicated three times. The study was a factorial type fitted in a Randomized Complete Block Design (RCBD). Phenological data collected were number of days to 50% tasselling, anthesis, silking and physiological maturity during the two cropping seasons. The study showed highly significant ( $P \leq 0.01$ ) decreases in days to 50% tasselling, anthesis, silking and maturity for the 2000 and 2001 cropping seasons with the sole application of 60kgP<sup>-1</sup>. Furthermore, the mean performance of phenological parameters showed the shortest days to silking with the combined application of 120kgNha<sup>-1</sup> and 60kgP<sup>-1</sup>, indicating that P application is critical for maize to show early phenological traits.

Corresponding Author's E-mail Address:  
e-mail: [ekwaghe16@gmail.com](mailto:ekwaghe16@gmail.com)

ISSN– Online 2736-1411

Print 2736-142X

© Publishing Realtime. All rights reserved.

### 1.0 Introduction

Maize ranks third in world's cereals after wheat and rice (Imran 2015). It is a short duration crop capable of producing large quantity of food grain. Among all the crop plants, maize is the most versatile one as it has great nutritive value; 72% starch, 10% protein, 4.8% 8.5% fibre, 3 % sugar and 17% ash (Chaudery, 1983). Of the three major developing regions of the world, Africa is the only one in which the index per capita food production has declined steadily during the last decade. Unless the present trend is reversed, 25 to 30 years from now, Africa will have the

world's largest deficit in cereals both in absolute and relative terms the challenges facing African agriculture in the next 20 years are immense, as the prospects of importing food to offset these deficits are not economically feasible.

To meet these challenges will require drastic increases in productivity of the principal staples of the continent through any well-articulated research and development. Maize more than any crop offers the promise of meeting Africa's food needs as no other cereals can be used in many ways as maize (RMRDC, 2004). Maize (*Zea mays*) is a member of the grass family poacea to which all cereals

belong. Also called corn, in the USA, is a monoceous plant which bears female organs, the pistil in separate flowers and same plant. The male inflorescence is commonly referred to as tassel occupying the terminal position on the main axis (Kochler, 1986). The silk of the female flower develops from the growing point of the flower. It elongates through the length of the husks propelled by the growth of an intercalary main stem located at the base. Each silk continues to grow until it is pollinated and fertilization takes place (Duncan, 1975).

Seasonal fluctuations in annual rainfall has made it difficult to achieve optimum yield of annual crops such as maize, either due to poor germination when rain ceases for some time after the initial onset or when the rain ceases before the completion of vegetation growth. However, the seed of improved variety and fertilizer are the main factors in enhancing the output of maize (Dowswell, et al., 1996). The soils in the study area are deficient in phosphorus and are alkaline, thus such soils have higher phosphorus fixation capacity and therefore higher demand for phosphorus (Hussein and Haqq, 2000). According to (Onasanya et al., 2009), appropriate type of fertilizer can increase the yield by 50 %. Phosphorus deficiency, therefore, is invariably a common crop growth and yield limiting factor in unfertilized especially in soils high in calcium carbonate which reduces p solubility (Ibrieki et al. 2005)

Phosphorus plays an indispensable role as a universal fuel for all biochemical activity in living cells and also plays a significant role in the breakdown of carbohydrates or other food products by photosynthesis in plants, which is usually supplied in the form of ATP. Other important role of phosphorus is in the stimulation of early maturity (Miller and Donahue, 1990). Mallarino et al. (1999) confirmed this when they reported that soil phosphorus availability during maize seedling development is critical for early growth and development of maize. Phosphorus can significantly increase vegetative growth and grain yield (Reghurum et al., 2000) as it plays many roles in many physiological processes that occur within a developing and maturing plant and is involved in enzymatic reactions in the plant.

According to Ibrahim and Kandi, (2007) and Amnanullah et al., (2009a), large quantities of phosphorus is found in seed, and it is considered essential for seed formation, inflorescence, grain formation, ripening and reproductive parts of maize plant, therefore, phosphorus in adequate amount is necessary for earlier maturity, rapid growth and improves the quality of vegetative growth and grain development in maize production. Phosphorus is essential in cell division because it is a constituent of element of nucleoproteins which are involved in the cell reproduction process (Imran 2015).

The objectives of this study were to evaluate the effects of phosphorus (P) on phenological traits of maize (*zea mays* L.) in Yola, North-Eastern Nigeria, and to make recommendations based on the results of the study to farmers in Yola and environs.

## 2.0 Materials and Methods

### 2.1 Description of the study area

The field experiments were located at the Research and Teaching Farm of the Department of Crop Production and Horticulture, Faculty of Agriculture, Federal University of Technology Yola. Yola located in the Northern Guinea Savannah ecological zone of Nigeria within latitude  $9^{\circ}19'N$  and longitude  $12^{\circ}30'E$  at an altitude of 185.9m above sea levels (Bashir, 2000). The area had a total rainfall of

745.5 mm and 690 mm in 2000 and 2001 cropping season. A total of 745.5 mm and 618.4 mm of rainfall were recorded during the crop life cycle of maize in 2000 and 2001 of rainfall were respectively (Table 1a and 1b). Rainfall was distributed over 150 and 160 days and can effectively support high maize production. Yola and its environment have high solar radiation which indicated high photosynthetic potentials for crops like maize that are C4 and require warm temperature.

### 2.2 Experiment design and treatment

The experimental design was a randomized complete block design (RCBD) with three replications during both the 2000 and 2001 cropping seasons. The treatments were 4 levels of nitrogen, and 2 levels of phosphorus which were factorially arranged resulting in 24 treatments. The levels of nitrogen (hereafter designated as N) were 0, 60, 120 and 180 kgN ha<sup>-1</sup> and phosphorus (hereafter designated as P) levels were 0 and 60 kgP<sup>-1</sup>. The total land area from the experiment was 430.5 m<sup>2</sup> which were demarcated and labelled into small plots of 10.5m<sup>2</sup> each made up of 3m x 3.5m. Each plot was demarcated into 5 rows with each row consisting of 12 hills. Composite soil samples were taken from each plot bulked, labelled, air dried in the shade and was taken to the laboratory to analyse for physicochemical properties. Particle size distribution was determined following the Walkey and Black method of chromic acid wet oxidation method (Nelson and Sommers, 1996). Total N was determined by the macro-kjedal digestion method (Bremner, 1996); available P was determined using the bicarbonate method (Olsen et al. 1985). Soil exchangeable cations (Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup>), were extracted with 1N NH<sub>4</sub> Acetate solution, but while Na<sup>+</sup> and K<sup>+</sup> were determined using the flame photometer, Ca<sup>2+</sup> and Mg<sup>2+</sup> were determined by titration with 0.02 N EDTA solution (Page et al. 1982).

### 2.3 Agronomic Practices

A popular maize hybrid variety, New Kaduna White sourced from Premier Seeds Kaduna was used for the experiment. Planting was done in early July for the two cropping seasons. The spacing between rows was 75 cm while the spacing within rows was 25 cm which gave a plant population of 53,333.33 plants per hectare. Basal application of 60kgP<sup>-1</sup> was applied at planting in all the plots except for the control where no P was applied. N fertilizer was applied in two split doses. 30 kgP<sup>-1</sup> at planting while the remaining 30 kgP<sup>-1</sup> was applied four weeks later. Weed control was done using a mixture of Atrazine and Garamazone herbicides which was supplemented with manual weeding as the field was removed.

### 2.4 Data Collection

Data on soil nutrient status, growth and phenological parameters were collected. Phenological parameters measured include days to 50% tasselling, anthesis, silking and maturity. All data were subjected to analysis of variance (ANOVA) using Statistical Analysis Software (SAS, 2002). The difference among significant treatment means were tested using least significant difference (LSD) at 5% level of significance.

## 3.0 results and Discussions

The meteorological information of the trial site for the 2000 and 2001 season is given in Tables 1a and 1b. Allaby (1998) reported that the factors of weather that affect growth and development of maize are temperature, rainfall, amount of sunshine and relative humidity. The total rainfall in Yola during the 2000 cropping season was 775.45 mm as against 690.00 mm for the 2001 cropping season. This was considered sufficient for optimum maize

production. It is mostly reported that maize is sensitive to drought stress during pollination when delayed emergence of silk may reduce fertilization (Jones, 1985). Drought stress as late as two to three weeks following 50% silking may also seed number (Frey, 1981). Similarly, temperature which is very important for phenological development and other physiological processes during growth and development of the crop were found to vary during the 2000 and 2001 cropping seasons. The maximum temperature during the month of June, July

and August, which was 38.8°C for year 2000 and 35.9°C for year 2001 was sufficient to hasten flowering and ear development.

The physical and chemical properties of the experimental sites are presented in Table 2. The results showed that the soils ranged from slightly acidic to moderately acidic sandy loam with an average of 67% sand, 20% silt and 13% clay. The soils were therefore generally coarse textured and well drained, characteristically low in org. C (0.74%), Avail. P (1.25 mgkg<sup>-1</sup>), exchangeable bases (0.61 cmolk<sup>-1</sup>) and CEC (2.81 cmolk<sup>-1</sup>).

Table 1a: Weather information during the 2000 cropping season in Federal University of Technology, Yola

| Month     | SRAD (hrs/day) | T MAX °C | T MIN °C | RAINFALL (mm) |
|-----------|----------------|----------|----------|---------------|
| Jan       | 5.6            | 55.6     | 21.3     | 0.00          |
| Feb       | 6.6            | 34.3     | 21.1     | 0.00          |
| March     | 11.1           | 40.6     | 25.3     | 0.00          |
| April     | 9.5            | 41.1     | 23.8     | 4.75          |
| May       | 7.5            | 37.8     | 26.1     | 99.9          |
| June      | 8.2            | 31.9     | 24.4     | 115.1         |
| July      | 8.1            | 31.0     | 23.7     | 122.7         |
| August    | 6.1            | 38.8     | 23.0     | 248.4         |
| September | 6.6            | 31.8     | 23.1     | 155.1         |
| October   | 7.4            | 34.6     | 23.4     | 29.5          |
| November  | 7.8            | 38.0     | 20.2     | 0.00          |
| December  | 5.1            | 34.5     | 18.3     | 0.00          |
| TOTAL     | 89.3           | 430      | 273.8    | 775.45        |

SRAD = Solar radiation; T MAX °C = Maximum Temperature; T MIN °C = Minimum Temperature

Table 1b: Weather information during the 2001 cropping season at Federal University of Technology Yola

| Month     | SRAD (hrs/day) | T MAX °C | T MIN °C | RAINFALL (mm) |
|-----------|----------------|----------|----------|---------------|
| Jan       | 6.4            | 35.3     | 19.0     | 0.00          |
| Feb       | 7.0            | 36.3     | 22.0     | 0.00          |
| March     | 10.2           | 41.5     | 26.1     | 0.00          |
| April     | 8.2            | 40.1     | 27.3     | 57.4          |
| May       | 8.9            | 38.9     | 26.8     | 122.2         |
| June      | 7.9            | 35.9     | 26.5     | 103.1         |
| July      | 6.9            | 31.0     | 23.8     | 155.2         |
| August    | 6.0            | 29.8     | 22.4     | 106.7         |
| September | 7.2            | 31.3     | 22.4     | 131.2         |
| October   | 18.2           | 31.6     | 20.0     | 14.2          |
| November  | 7.7            | 36.0     | 19.5     | 0.00          |
| December  | 5.8            | 35.0     | 19.5     | 0.00          |
| TOTAL     | 90.4           | 422.7    | 237      | 690           |

SRAD = Solar radiation; T MAX °C = Maximum Temperature; T MIN °C = Minimum Temperature

Table 2: Physicochemical properties of the soils of the experimental site for the 2000 and 2001 cropping seasons

| Soil characteristics                                  | Values for year 2000 | Values for year 2001 |
|-------------------------------------------------------|----------------------|----------------------|
| Sand (%)                                              | 65                   | 69                   |
| Silt (%)                                              | 21                   | 19                   |
| Clay (%)                                              | 14                   | 12                   |
| Soil texture                                          | Sandy loam           | sandy loam           |
| Soil pH in H <sub>2</sub> O                           | 6.27                 | 6.38                 |
| Organic carbon (%)                                    | 0.54                 | 0.40                 |
| Total N (%)                                           | 0.06                 | 0.08                 |
| Available P (mgkg <sup>-1</sup> )                     | 0.8                  | 0.9                  |
| Exchangeable Na <sup>+</sup> (cmolkg <sup>-1</sup> )  | 0.28                 | 0.32                 |
| Exchangeable K <sup>+</sup> (cmolkg <sup>-1</sup> )   | 0.28                 | 0.37                 |
| Exchangeable Mg <sup>2+</sup> (cmolkg <sup>-1</sup> ) | 0.41                 | 0.62                 |
| Exchangeable Ca <sup>2+</sup> (cmolkg <sup>-1</sup> ) | 0.33                 | 0.46                 |
| CEC (cmolkg <sup>-1</sup> )                           | 2.76                 | 2.85                 |

CEC = Cation exchange capacity; P =phosphorus; Na = Sodium; K = Potassium; Mg = Magnesium; Ca = Calcium

Source: Soil Science Laboratory, Federal University of Technology, Yola

During the 2000 cropping season, highly significant differences among means ( $P \leq 0.01$ ) were observed due to P treatments for days to 50% tasseling, anthesis, and maturity. However, days to 50 % silking showed significant differences among means ( $P \leq 0.05$ ), due to P treatment during the same cropping season (Table 3a). During the 2001 cropping season, highly significant differences among means ( $P \leq 0.01$ ) were observed for all the phenological traits measured (days to 50% tasseling, anthesis, silking and maturity) due to P treatments (Table 3b).

Furthermore, for both the 2000 and 2001 cropping seasons, the mean performances of all the phenological traits observed

showed significant reduction in days to 50% tasselling, anthesis, silking and physiological maturity (Tables 4a and 4b). These significant reduction in days observed in all the phenological traits for the two cropping seasons could be attributed to the indispensable role played by P in its many compounds in cell stimulation and hastening of maturity. This is consistent with the reports by Miller and Donahue (1990) and Mallarino et al., (1999) who reported that P plays an indispensable role as a universal fuel for all biological activity in living cells, which is critical to early growth and development of maize.

Table 3a: Analysis of variance Table showing the mean squares of phonological parameters of maize measured during the 2000 cropping seasons

| Source of variations | d.f. | Days to 50% tasselling | Days to 50% anthesis  | Days to 50% silking | Days to 50% maturity |
|----------------------|------|------------------------|-----------------------|---------------------|----------------------|
| Replication          | 3    | 0.6667                 | 0.1667                | 2.042               | 4.29                 |
| N                    | 3    | 0.5000 <sup>ns</sup>   | 1.8194 <sup>**</sup>  | 1.042 <sup>ns</sup> | 3.92 <sup>ns</sup>   |
| P                    | 1    | 20.16678 <sup>**</sup> | 18.3750 <sup>**</sup> | 9.375 <sup>*</sup>  | 210.04 <sup>**</sup> |
| N x P                | 3    | 0.9444 <sup>ns</sup>   | 2.2639 <sup>**</sup>  | 4.153 <sup>*</sup>  | 15.15 <sup>ns</sup>  |
| Error                | 14   | 0.7143                 | 0.3571                | 1.280               | 13.29                |

\*\* = Significantly different at ( $P \leq 0.01$ ); \* = Significantly different at ( $P \leq 0.05$ );

ns = Not significantly different at ( $P \leq 0.05$ ); d.f. = degrees of freedom

3b: Analysis of variance Table showing the mean squares of phenological parameters of maize measured during the 2001 cropping seasons

| Source of variations | d.f. | Days to 50% tasselling | Days to 50% anthesis | Days to 50% silking  | Days to 50% maturity  |
|----------------------|------|------------------------|----------------------|----------------------|-----------------------|
| Replication          | 3    | 2.375                  | 0.365                | 1.781                | 10.781                |
| N                    | 3    | 2.875 <sup>ns</sup>    | 2.031 <sup>ns</sup>  | 3.531 <sup>ns</sup>  | 7.365 <sup>*</sup>    |
| P                    | 1    | 36.125 <sup>**</sup>   | 47.531 <sup>**</sup> | 47.431 <sup>**</sup> | 132.031 <sup>**</sup> |
| N x P                | 3    | 1.708 <sup>ns</sup>    | 1.365 <sup>ns</sup>  | 1.781 <sup>ns</sup>  | 2.865 <sup>ns</sup>   |
| Error                | 21   | 1.042                  | 1.103                | 1.353                | 2.424                 |

\*\* = Significantly different at ( $P \leq 0.01$ ); \* = Significantly different at ( $P \leq 0.05$ );

ns = Not significantly different at ( $P \leq 0.05$ ); d.f. = degrees of freedom

Table 4a: Mean performance of phenological parameters of maize due to P treatments measured during the 2000 cropping season

| P rates (kg ha <sup>-1</sup> ) | Days to 50% tasselling | Days to 50% anthesis | Days to 50% silking | Days to 50% maturity |
|--------------------------------|------------------------|----------------------|---------------------|----------------------|
| 0                              | 63.33                  | 65.08                | 65.58               | 83.42                |
| 60                             | 61.50                  | 63.33                | 65.33               | 77.50                |
| LSD                            | 0.690                  | 0.523                | 0.991               | 3.192                |
| F Prob.                        | 0.001                  | < 0.001              | 0.017               | < 0.001              |

LSD = Least Significant Difference

Table 4b: Mean performance of phenological parameters of maize due to P treatments measured during the 2001 cropping season

| P rates (kg ha <sup>-1</sup> ) | Days to 50% tasselling | Days to 50% anthesis | Days to 50% silking | Days to 50% maturity |
|--------------------------------|------------------------|----------------------|---------------------|----------------------|
| 0                              | 53.75                  | 56.69                | 60.56               | 89.50                |
| 60                             | 51.63                  | 51.25                | 58.13               | 85.44                |
| LSD                            | 1.061                  | 0.772                | 0.855               | 1.145                |
| F Prob.                        | < 0.001                | < 0.001              | < 0.001             | < 0.001              |

LSD = Least Significant Difference

#### 4.0 Conclusion

From the results of this study, it can be concluded that the basal application of 60kg P ha<sup>-1</sup> achieved significant reduction in days to 50% tasselling, anthesis, silking and physiological maturity of maize in Yola and its environs, indicating that P application is critical for maize to show early phenological traits. It is also recommended that the experiment be

conducted across different locations with varied ecology in Nigeria. Furthermore, similar experiment should be conducted with other varieties of maize in the same ecological zone to ascertain the effects of P on phenological traits of other varieties.

## References

- Allaby, M. (1988). *Macmillan Dictionary of Environment*, 3<sup>rd</sup> Edition. London. Pp. 358-359
- Amanullah, R/A., Khattak, S., Khali, K.(2009b). Effects of plant density and N on phenology and yield of maize. *J. Plant Nutr.* 32 (3): 245-259). .
- Bashir, A. (2000). Water quality and outbreak of diseases in Yola area of Adamawa State. An M.Sc. dissertation submitted to the Department of Geography, Federal University of Technology, Yola.
- Black, C.A. (1965). Method of Soil Analysis. *Agronomy* No. 9 part 2. Am Soc. Agron. Madison Wisconsin
- Bremmer, J.M. (1996). Total Nitrogen. In Black C.A. (ed). *Method of Soil Analysis*, part 2. ASA, Madison, Wisconsin. Vol. 9.pp. 914-926
- Chaundhary, A. R. (1983). Maize in Pakistan. Punjab Agricultural Research Coordination Board, University of Agriculture,
- Downswell, C.R., Paltiwai, R.F. and Cantrell, R.L. (1996). *Maize in the third world*. West View Press. pp. 268
- Duncan, W.G. (1975). Maize Crop physiology. Evans (ed). London: Cambridge University Press, pp. 23-50.
- Frey, N. M. (1981). Dry matter accumulations in maize kernels of maize. *Crop Science* Vol. 1: pp 118-122
- Hussein, M.Z. and Haqq, I.U. (2000). Phosphorus absorption capacities of KPK soils. In: Proceedings of Symposium on integrated plant nutrient management held at Islamabad on 8-10 November. 1999. Pp. 284-296
- Ibrahim, S. A. and Kandil, H. (2007). Growth yield and chemical constituents of Corn (*Zea mays*) affected by nitrogen and phosphorus fertilization under different irrigation Intervals. *Journal of Applied Sciences Research*, Vol. 3, pp 1112-11120
- Ibrieki, H.J., Ryan, A.C., Ulger, G... Buyuk, B., Cakir, K., Korkmaz, E., Karnez, G., Ozgenturk, O. and Konuskan, C. (2005). Maintenance of P fertilizer and residual P effect on corn production. *Nigerian J. Soil Sci.* Vol 2.pp. 11271279-1286
- Imran, I. (2015). Effect of germination on proximate composition of two maize cultivars. *J. Bio and H. Care*. Vol.5 (3), pp 123- 128
- Jones, C.A. (1985). *C<sub>4</sub> Grasses and Cereals. Growth, Development and stress response*. New York: Wiley, pp.16
- Kochler, S.L. (1986). *Tropical crops. A text book of economic botany*. London: Macmillan pp. 88-98
- Malarrino, A.P.; Bordolli, J.M and Borgues, R. (1999). Phosphorus and potassium placement effects on early growth and nutrient uptake of No-Till Corn and relationships with grain yield. *Agronomy Journal*, Vol.91, pp7-45
- Miller, R. W. and Donahue, R. L. (1990). *An introduction to soil and plant growth*, 6<sup>th</sup> Edition. Prentice Hall. pp. 250
- Nelson, D.M. and Sommers, L. (1996). Total carbon, organic carbon and organic matter. In: D.L. Sparks (eds). *Methods of Soil Analysis part 3- chemical methods*. SSSA Book Series. Vol. 5, Madison Wisconsin, USA, pp. 961-1010
- Olsen, S.R.; Watanabe, F.S. and Dean, L.A. (1985). Estimation of available P in the soil by extraction with sodium bicarbonate. USDA: pp939
- Page, A. L.; Miller, P.H. and Keeney, D.R. (1982). *Methods of Soil Analysis Part 2. Chemical and Microbiological properties*, 2<sup>nd</sup> Edition. USA: ASA. Inc. Madison Wisconsin
- Reghurum, K., Singn, T.A., and Vasta, N.K. (2000). Effect of phosphate fertilization of different water solubility on maize and residual on wheat in a mollisol. *Pantnagar J. of Res.*, Vol. 29. pp. 40-43
- RMRDC. (1997). *Raw Materials Sourcing for Manufacturing in Nigeria*. 3<sup>rd</sup> Edition, Abuja, pp146.